

**Toxoplasma gondii, Strain SF47**

**Catalog No. NR-49196**

**Product Description:** *Toxoplasma gondii* (*T. gondii*), strain SF47 is a recombinant F1 clone selected from progeny of a genetic cross between a sinfungin-resistant line of the highly virulent Type I GT-1 strain (GT1-SNF<sup>R</sup>) and a 5-fluoro-2'-deoxyuridine-resistant line of the non-virulent Type II ME49 strain (ME49 FUDR<sup>R</sup>).

**Lot<sup>1,2</sup>: 64223570**

**Manufacturing Date: 06MAY2016**

TEST	SPECIFICATIONS	RESULTS
<b>Cell Morphology</b>	Report results	Refractile and oval-shaped
<b>PCR Assay of Extracted DNA<sup>3,4</sup></b> AK56 locus	~ 520 base pair amplicon	~ 520 base pair amplicon
<b>Genotypic Analysis<sup>3,4</sup></b> Sequencing of AK56 locus (~ 510 base pairs) AK56 locus ( <i>MfeI</i> digestion)	Consistent with <i>T. gondii</i> Consistent with parental Type I strain	Consistent with <i>T. gondii</i> (Figure 1) Consistent with parental Type I strain
<b>Viable Cell Count by Hemacytometry (pre-freeze)</b>	> 10 <sup>6</sup> cells/mL	2.7 × 10 <sup>7</sup> cells/mL
<b>Viability (post-freeze)<sup>5</sup></b>	Viable parasites	Viable parasites
<b>Sterility (21-day incubation)</b> Harpo's HTYE broth <sup>6</sup> , 37°C and 26°C, aerobic Tryptic Soy broth, 37°C and 26°C, aerobic Sabouraud Dextrose broth, 37°C and 26°C, aerobic DMEM with 10% FBS, 37°C, aerobic Sheep Blood agar, 37°C, aerobic Sheep Blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
<b>Mycoplasma Contamination</b> DNA Detection by PCR	None detected	None detected

<sup>1</sup>NR-49196 was produced by cultivation of the deposited material in human foreskin fibroblast cells (ATCC<sup>®</sup> CRL-1634<sup>™</sup>) with cell cultivation medium for parasites (ATCC<sup>®</sup> medium 2222: adjusted to contain 10% heat-inactivated fetal bovine serum). The culture was propagated for 4 days at 37°C in an aerobic atmosphere with 5% CO<sub>2</sub> until lysis of the host cell monolayer was reached.

<sup>2</sup>Quality control testing completed on post-freeze material unless specified as pre-freeze.

<sup>3</sup>PCR amplification of the AK56 locus was performed. Samples were subjected to restriction enzyme digestion typing by agarose gel electrophoresis.

<sup>4</sup>Primer sequences, annealing temperatures, and conditions for restriction enzyme digestion may be obtained at the *Toxoplasma* Genome Map website ([Toxoplasma Genome Map](http://ToxoplasmaGenomeMap.org)).

<sup>5</sup>Viable cells and signs of infection were seen after 4 days under cultivation conditions at 37°C.

<sup>6</sup>Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

**Figure 1: AK56 (Chromosome II) Amplicon Sequence**

```

GTCCTTTTCC CCACTGCTTT TATTAGGTTT TTCCGTGTTT TCGCGGAGTC GTCTGAGCTC GGCAC'TCGCT GCTTTCCAAA
ATCTCGTTTC AACGTATCGC GGCGCCGTCA CCGCGCGCAA TCCACTGTGA TGCATGATTC TGTTTCTAAA AACTGCGCAT
TTTAGCCGGC TCGTTTTTGC ATACGTTTGG ACCATAAAAC CTCGTATTGT TGAAGAAGAA TGCAATTGGT GTCTGTGCTG
ATCACCGTAT GAAAATCGGC GTGTCTCGCC CCCTGCCGTG TCGCGTCCG CTTTTTGCGA CCCC'GGTACA CCCGTTTTTT
GTGGTCAGCG AGGAACGCAC TTTTGCTGTT ATTGTTCACT TTTCAGCGTA AACTGACCC CTTTCATCGT GGCAGGAAAC
GA'ACTCTCAG CAAGAATTTT CGAGCACTAC TGC'GTGCAG CAGCCTAGTG GGGTGGACAC GCATGTGCAG GACGGACAGA
AACTGCAAGC TTGTTCCGCA GGCTAAA'ACT C
    
```

**Date:** 04 NOV 2016

**Signature:**



BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

