

## **Certificate of Analysis for NR-45992**

## Staphylococcus aureus, Strain C1999000193

## Catalog No. NR-45992

**Product Description**: *Staphylococcus aureus* (*S. aureus*), strain C1999000193 was isolated in 1999 from pleural fluid of a 13-year-old female with necrotizing pneumonia and severe sepsis in rural Minnesota, USA. *S. aureus*, strain C1999000193 is a community-associated methicillin-resistant *S. aureus* (CA-MRSA) strain.

Lot<sup>1</sup>: 2145 Manufacturing Date: 03NOV2016

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis		
Cellular morphology	Gram-positive cocci	Gram-positive cocci
Colony morphology <sup>2</sup>	Report results	Circular, convex, entire, smooth and
Colony morphology	Troport Todato	cream (Figure 1)
Motility (wet mount)	Report results	Non-motile
Hemolysis <sup>3</sup>	Report results	β-hemolytic
Biochemical characterization	Report results	p nemoryte
Catalase	Positive	Positive
Coagulase <sup>4</sup>	Report results	Positive
VITEK® 2 Compact (GP card)	≥ 90% probability of being <i>S. aureus</i>	S. aureus (99% probability) <sup>5</sup>
	2 90 % probability of being 3. aureus	3. aureus (99 % probability)
Antibiotic Susceptibility Profile VITEK® (AST-GP71 card)6		
Beta-lactamase <sup>7</sup>	Poport regulto	Positive
	Report results Report results	Positive
Cefoxitin screen		
Benzylpenicillin	Report results	Resistant (≥ 0.5 μg/mL)
Oxacillin	Resistant	Resistant (≥ 4 μg/mL)
Gentamicin	Sensitive	Sensitive (≤ 0.5 μg/mL)
Ciprofloxacin	Sensitive	Sensitive (≤ 0.5 μg/mL)
Levofloxacin	Report results	Sensitive (≤ 0.12 µg/mL)
Moxifloxacin	Report results	Sensitive (≤ 0.25 μg/mL)
Clindamycin (inducible resistance)	Report results	Negative
Erythromycin	Sensitive	Sensitive (≤ 0.25 μg/mL)
Clindamycin	Sensitive	Sensitive (≤ 0.25 µg/mL)
Quinupristin/dalfopristin	Sensitive	Sensitive (≤ 0.25 μg/mL)
Linezolid	Sensitive	Sensitive (= 2 µg/mL)
Daptomycin	Report results	Sensitive (= 0.25 µg/mL)
Vancomycin	Sensitive	Sensitive (= 1 µg/mL)
Minocycline	Report results	Sensitive (≤ 0.5 µg/mL)
Tetracycline	Report results	Sensitive (≤ 1 μg/mL)
Tigecycline	Report results	Sensitive (≤ 0.12 µg/mL) <sup>8</sup>
Nitrofurantoin	Report results	Sensitive (≤ 16 μg/mL)
Rifampicin	Report results	Sensitive (≤ 0.5 µg/mL)
Trimethoprim/sulfamethoxazole	Sensitive	Sensitive (≤ 10 μg/mL)
Etest® antibiotic test strips9		
Chloramphenicol <sup>10</sup>	Report results	Sensitive (= 4 µg/mL)
Teicoplanin <sup>10</sup>	Sensitive	Sensitive (= 1 µg/mL)
Genotypic Analysis		
Sequencing of 16S ribosomal RNA gene	≥ 99% sequence identity to	100% sequence identity to
(~ 1500 base pairs)	S. aureus type strain	S. aureus type strain
	(GenBank: L37597)	(GenBank: L37597)
	Consistent with expected colony	Consistent with expected colony
Purity (post-freeze) <sup>11</sup>	morphology	morphology
	Погрноюду	тюгрпоюду
Viability (post-freeze) <sup>2</sup>	Growth	Growth

BEI Resources www.beiresources.org E-mail: <a href="mailto:contact@beiresources.org">contact@beiresources.org</a>
Tel: 800-359-7370

Fax: 703-365-2898



SUPPORTING INFECTIOUS DISEASE RESEARCH

## Certificate of Analysis for NR-45992

10 average storie 04000000400 was demonstrad to DEL Description on next of the NADCA of

1S. aureus, strain C1999000193 was deposited to BEI Resources as part of the NARSA collection. NR-45992 was produced by inoculation of the deposited material into Tryptic Soy broth and grown 1 day at 37°C in an aerobic atmosphere. Broth inoculum was added to Tryptic Soy agar kolles which were grown 1 day at 37°C in an aerobic atmosphere to produce this lot.

<sup>2</sup>1 day at 37°C in an aerobic atmosphere on Tryptic Soy agar

<sup>3</sup>1 day at 37°C in an aerobic atmosphere on Tryptic Soy agar with 5% defibrinated sheep blood

<sup>4</sup>4 hours at 37°C in rabbit serum with 0.15% EDTA (Coagulase Plasma BBL™ 240827)

<sup>5</sup>Percent probabilities above 90% indicate a close match to the typical biochemical pattern for the given organism, with a percent probability of 99% being a perfect match between the test reaction pattern and the unique biochemical pattern of the given organism or organism group. For additional information, please refer to O'Hara, C.M. and J. M. Miller. "Evaluation of the VITEK 2 ID-GNB Assay for Identification of Members of the Family Enterobacteriaceae and Other Nonenteric Gram-Negative Bacilli and Comparison with the VITEK GNI+ Card." J. Clin. Microbiol. 41 (2003): 2096-2101. PubMed: 12734254.

<sup>6</sup>Minimum Inhibitory Concentration (MIC): MIC Interpretation Guideline: CLSI M100-S22 (2012)

<sup>7</sup>The production of beta-lactamase was detected using a Cefinase™ Paper Disc (BBL™ 231650).


<sup>8</sup>MIC Interpretation Guideline: EUCAST Version 4.0 (2014)

<sup>9</sup>1 day at 37°C in an aerobic atmosphere on Mueller Hinton agar

<sup>10</sup>For both chloramphenicol (bioMérieux Etest<sup>®</sup> 412308) and teicoplanin (bioMérieux Etest<sup>®</sup> 412459), a MIC ≤ 8 μg/mL is sensitive, a MIC = 16 μg/mL is intermediate and a MIC ≥ 32 μg/mL is resistant.

<sup>11</sup>Purity of this lot was assessed for 7 days at 37°C in an aerobic atmosphere with 5% CO<sub>2</sub> on Tryptic Soy agar with 5% defibrinated sheep blood.





**Date:** 13 DEC 2016

Signature:

**BEI Resources Authentication** 

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

BEI Resources www.beiresources.org E-mail: contact@beiresources.org

Tel: 800-359-7370 Fax: 703-365-2898