

Influenza B Virus, B/New York/1061/2004 (Yamagata Lineage)

Catalog No. NR-48663

Product Description: Cell lysate and supernatant from Madin-Darby Canine Kidney (MDCK) cells¹ infected with Influenza B Virus, B/New York/1061/2004 (Yamagata Lineage)

Passage History: RhMK1/C2 (Prior to deposit at BEI Resources/BEI Resources); RhMK# = Number of passages in primary Rhesus monkey kidney cells; V# = Number of passages in MDCK cells

Lot²: 64373901

Manufacturing Date: 24SEP2016

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity in MDCK Cells ¹	Cell rounding and detachment	Cell rounding and detachment
Sequencing of NB Protein and Neuraminidase Coding Region (835 nucleotides)	Consistent with B/New York/1061/2004 (Yamagata Lineage)	99% identity with B/New York/1061/2004 (Yamagata Lineage) (GenBank: CY174379)
Titer by TCID ₅₀ Assay ^{3,4} in MDCK cells ¹	Report results	1.6 × 10 ⁶ TCID ₅₀ per mL
Sterility (21-day incubation) Harpo's HTYE broth ⁵ , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Blood agar, 37°C, aerobic Blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO ₂	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA detection by PCR of extracted Test Article nucleic acid	None detected None detected	None detected None detected

¹MDCK; ATCC® CCL-34™

²Grown in Eagle's Minimum Essential Medium (ATCC® 30-2003) supplemented with 0.125% bovine serum albumin (Gibco 15260-037) and 1.0 µg/mL L-1-tosylamido-2-phenylethyl chloromethyl ketone (TPCK)-treated trypsin for 3 days at 33°C and 5% CO₂

³The Tissue Culture Infectious Dose 50% (TCID₅₀) endpoint is the 50% infectious endpoint in cell culture. The TCID₅₀ is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID₅₀ provides a measure of the infectious titer (or infectivity) of a virus preparation.

⁴7 days at 33°C and 5% CO₂

⁵Atlas, Ronald M. Handbook of Microbiological Media. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

Date: 26 APR 2017

Signature: 

BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

