

Cryptococcus gattii, Strain C17

Catalog No. NR-50431

Product Description: *Cryptococcus gattii* (*C. gattii*), strain C17 was isolated from a human bronchial wash in the Pacific Northwest region of North America. *C. gattii*, strain C17 was deposited as lineage VGI and resistant to azoles.

Lot¹: 2208

Manufacturing Date: 19DEC2016

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis Cellular morphology ² Colony morphology ² Canavanine-glycine-bromthymol blue (CGB) differential medium ³	Report results Report results Report results	Globose or subglobose; single or budding (Figure 1A) Entire, smooth, mucoid and cream (Figure 1B) Blue
Genotypic Analysis Sequencing of partial 18S rRNA gene, internal transcribed spacer (ITS) 1, 5.8S rRNA gene, ITS 2, partial 28S rRNA (~ 520 base pairs) Sequencing of 26S rRNA gene (~ 620 base pairs)	≥ 99% sequence identity to <i>C. gattii</i> (GenBank: FJ914888.1) ≥ 99% sequence identity to <i>C. gattii</i> (GenBank: KC171326.1)	99.2% sequence identity to <i>C. gattii</i> (GenBank: FJ914888.1) 99.8% sequence identity to <i>C. gattii</i> (GenBank: KC171326.1)
Antimicrobial Susceptibility^{4,5} Fluconazole	Report MIC (µg/mL)	0.19 µg/mL to 0.25 µg/mL
Purity⁶ Nutrient broth with 0.1% Yeast Extract at 25°C Nutrient broth with 0.1% Yeast Extract at 37°C	No bacterial growth No bacterial growth	No bacterial growth No bacterial growth
Viability (post-freeze)²	Growth	Growth

¹NR-50431 was produced by inoculation of the deposited material onto Yeast Mold agar and incubated for 5 days at 25°C in an aerobic atmosphere. Cells were harvested from agar with 20% glycerol.

²3 days at 25°C in an aerobic atmosphere on modified Sabouraud Dextrose agar


³2 days at 26°C in an aerobic atmosphere. CGB medium differentiates *C. gattii* from *C. neoformans* based on the ability of *C. gattii* isolates to grow in the presence of L-canavanine and to assimilate glycine as a sole carbon source, resulting in a blue color. *C. neoformans* isolates will show yellow to light green on CGB medium [McTaggart, L., et al. "Rapid Identification of *Cryptococcus neoformans* var. *grubii*, *C. neoformans* var. *neoformans*, and *C. gattii* by Use of Rapid Biochemical Tests, Differential Media, and DNA Sequencing." *J. Clin. Microbiol.* 49 (2011): 2522-2527. PubMed: 21593254.].

⁴2 days at 35°C in an aerobic atmosphere on RPMI 1640 agar with MOPS and 2% glucose (Remel R04067)

⁵bioMérieux Etest® 510858

⁶Clarity of broth was determined by visual inspection after 5 days in an aerobic atmosphere.

Figure 1: Cellular and Colony Morphologies



Certificate of Analysis for NR-50431

Date: 12 JUN 2017

Signature: 

BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

