

Certificate of Analysis for NR-48749

Mycobacterium tuberculosis, Strain 11511-0

Catalog No. NR-48749

www.beiresources.org

Product Description: *Mycobacterium tuberculosis* (*M. tuberculosis*), strain 11511-0 was isolated in October 2012 from a subculture of a strain originally isolated from a patient with pulmonary tuberculosis in the Republic of South Africa. *M. tuberculosis*, strain 11511-0 was deposited as a multi-drug resistant (MDR) strain with resistance to isoniazid and rifampin.

Lot¹: 63950966 Manufacturing Date: 08JAN2016

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis ²		
Cellular morphology	Gram-positive rods	Gram-positive rods
Colony morphology ³	Report results	Irregular, raised, entire, rough and cream (Figure 1)
Growth rate	≥ 7 days	22 days
Growth at 26°C	Negative	Negative
Growth at 37°C	Positive	Positive
Acid-fast stain	Positive (red colonies)	Positive (red colonies)
Pigmentation in the dark (Scotochromogen)	Negative (no pigment)	Negative (no pigment)
Photoinduction for 1 hour (Photochromogen)	Negative (no pigment)	Negative (no pigment)
Nonchromogen (no pigment)	Positive (no pigment)	Positive (no pigment)
Biochemical tests	l contro (no pigmont)	1 collive (ne pigmont)
Niacin production ⁴	Positive	Positive
Nitrate reduction	Positive	Positive
Pyrazinamidase	Positive	Positive
	1 001410	1 Goldvo
Antibiotic Susceptibility Profile		
Sensititre TM System ^{5,6}	Daniel and an action	0.05/1
Amikacin	Report results	0.25 μg/mL
Cycloserine	Report results	≤ 2 μg/mL
Ethambutol	Report results	4 μg/mL ⁷
Ethionamide	Report results	2.5 μg/mL ⁷
Isoniazid	Report results	2 μg/mL
Kanamycin	Report results	1.2 μg/mL
Moxifloxacin	Report results	0.5 μg/mL
Ofloxacin	Report results	1 μg/mL
Para-aminosalicylic acid	Report results	≤ 0.5 µg/mL ⁷
Rifabutin	Report results	> 16 µg/mL ⁷
Rifampin	Report results	> 16 µg/mL
Streptomycin	Report results	2 μg/mL ⁷
Genotypic Analysis		
Sequencing of Heat Shock Protein 65 gene	≥ 99% sequence identity to	100% sequence identity to
(~ 430 base pairs)	M. tuberculosis type strain	M. tuberculosis type strain
,	(GenBank: AL123456)	(GenBank: AL123456) ⁸
Purity (post-freeze)		
Middlebrook 7H10 agar with OADC enrichment ⁹	Growth consistent with expected	Growth consistent with expected
wild debiook /1110 agai with OADO enitchments	colony morphology	colony morphology
Tryptic Soy agar ⁹	Report results	No growth
Tryphic Goy again	Treport results	140 growth
Viability (post-freeze) ³	Growth	Growth

¹NR-48749 was produced by inoculation of the deposited material into Middlebrook 7H9 broth with ADC enrichment. Broth inoculum was added to Middlebrook 7H10 agar with OADC enrichment kolles, which were grown for 22 days at 37°C in an aerobic atmosphere with 5% CO₂ to produce this lot.

BEI Resources E-mail: contact@beiresources.org

Tel: 800-359-7370 Fax: 703-365-2898

²Information on Mycobacterium testing is available from Ribón, W. "Biochemical Isolation and Identification of Mycobacteria." <u>Biochemical Testing.</u> (2012) Jose C. Jimenez-Lopez (Ed.), InTech, http://www.intechopen.com/books/biochemical-testing/biochemical-isolation-and-identification-of-the-">http://www.intechopen.com/books/biochemical-testing/biochemical-isolation-and-identification-of-the-">http://www.intechopen.com/books/biochemical-testing/biochemical-isolation-and-identification-of-the-">http://www.intechopen.com/books/biochemical-testing/biochemical-isolation-and-identification-of-the-">http://www.intechopen.com/books/biochemical-testing/biochemical-isolation-and-identification-of-the-">http://www.intechopen.com/books/biochemical-testing/biochemical-isolation-and-identification-of-the-">http://www.intechopen.com/books/biochemical-testing/biochemical-isolation-and-identification-of-the-">http://www.intechopen.com/books/biochemical-testing/biochemical-isolation-and-identification-of-the-">http://www.intechopen.com/books/biochemical-testing/biochemical-isolation-and-identification-of-the-">http://www.intechopen.com/books/biochemical-testing/biochemical-isolation-and-identification-of-the-">http://www.intechopen.com/books/biochemical-testing/biochemical-isolation-and-identification-of-the-">http://www.intechopen.com/books/biochemical-testing/biochemical-isolation-and-identification-and-identifi



Certificate of Analysis for NR-48749


mycobacteria and Lévy-Frébault, V. V. and F. Portaels. "Proposed Minimal Standards for the Genus Mycobacterium and for Description of New Slowly Growing *Mycobacterium* Species." <u>Int. J. Syst. Bacteriol.</u> 42 (1992): 315-323. PubMed: 1581193. ³22 days at 37°C in an aerobic atmosphere with 5% CO₂ on Middlebrook 7H10 agar with OADC enrichment

⁴All mycobacteria produce niacin but only M. tuberculosis accumulates it, resulting in a positive test for M. tuberculosis.

⁵Sensititre[™] System *Mycobacterium tuberculosis* MIC Plate, Thermo Scientific[™], catalog number MYCOTB

⁸Also consistent with M. africanum, M. bovis, M. canettii, M. caprae and M. microti

Figure 1: Colony Morphology



Date: 16 JUN 2017

Signature:

BEI Resources Authentication

ATCC[®], on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

BEI Resources

www.beiresources.org

E-mail: contact@beiresources.org

Tel: 800-359-7370 Fax: 703-365-2898

⁶Minimum Inhibitory Concentration (MIC); No Clinical & Laboratory Standards Institute (CLSI) interpretations of the Sensititre™ System data for M. tuberculosis are currently available.

⁷For streptomycin, ethionamide, para-aminosalicylic acid, rifabutin and ethambutol, the endpoint for these drugs is determined by the well with approximately 80% inhibition of growth compared to the positive control well with no drug.

⁹Purity of this lot was assessed for 22 days at 37°C in an aerobic atmosphere with 5% CO₂.