

Mycobacterium florentinum*, Strain FI-93171T*Catalog No. NR-49073**

Product Description: *Mycobacterium florentinum* (*M. florentinum*), strain FI-93171T was isolated in 1993 from an excised lymph node of a 6-year-old girl in Italy.

Lot¹: 64362416**Manufacturing Date:** 04AUG2016

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis^{2,3}		
Cellular morphology	Rods	Rods
Colony morphology ⁴	Report results	Circular, low convex, entire, smooth and cream (Figure 1)
Growth rate	≥ 7 days	14 days
Growth at 45°C	Negative	Negative
Growth at 55°C	Report results	Negative
Acid-fast stain	Positive (red colonies)	Positive (red colonies)
Pigmentation in the dark (Scotochromogen)	Negative (no pigment)	Negative (no pigment)
Photoinduction for 1 hour (Photochromogen)	Negative (no pigment)	Negative (no pigment)
Nonchromogen (no pigment)	Positive	Positive
Biochemical tests		
Catalase	Report results	Positive
Catalase (semiquantitative)	Report results	Negative
Catalase (68°C)	Report results	Positive
Iron uptake	Report results	Negative
Nitrate reduction	Report results	Negative
Tween 80 hydrolysis	Negative	Negative
Urease	Positive	Positive
Growth in the presence of 5% sodium chloride	Negative	Positive ⁵
Growth in the presence of thiophene-2-carboxylic acid hydrazide (TCH)	Report results	Negative
Genotypic Analysis		
Sequencing of 16S ribosomal RNA gene (~1790 base pairs)	≥ 99% sequence identity to <i>M. florentinum</i> type strain (GenBank: AJ616230)	100% sequence identity to <i>M. florentinum</i> type strain (GenBank: AJ616230)
Digital DNA-DNA hybridization (dDDH) ⁶	≥ 70% for species identification	Not determined ^{7,8} (Table 1)
Purity (post-freeze)		
Middlebrook 7H10 agar with OADC enrichment ⁹	Growth consistent with expected colony morphology	Growth consistent with expected colony morphology
Tryptic Soy agar ⁹	Report results	Growth consistent with expected colony morphology
Viability (post-freeze)⁴	Growth	Growth

¹NR-49073 was produced by inoculation of the deposited material into Middlebrook 7H9 broth with ADC enrichment and grown for 15 days at 37°C in an aerobic atmosphere with 5% CO₂. Broth inoculum was added to Middlebrook 7H10 agar with OADC enrichment kolles, which were grown for 9 days at 37°C in an aerobic atmosphere with 5% CO₂ to produce this lot.

²Information on Mycobacterium testing is available from Ribón, W. "Biochemical Isolation and Identification of Mycobacteria." *Biochemical Testing*. (2012) Jose C. Jimenez-Lopez (Ed.), InTech, <http://www.intechopen.com/books/biochemical-testing/biochemical-isolation-and-identification-of-mycobacteria> and Lévy-Frébault, V. V. and F. Portaels. "Proposed Minimal Standards for the Genus *Mycobacterium* and for Description of New Slowly Growing *Mycobacterium* Species." *Int. J. Syst. Bacteriol.* 42 (1992): 315-323. PubMed: 1581193.

³Phenotypic characterization of *M. florentinum* was performed following Tortoli, E., et al. "*Mycobacterium florentinum* sp. nov., Isolated from Humans." *Int. J. Syst. Evol. Microbiol.* 55 (2005): 1101-1106. PubMed: 15879240.

⁴14 days at 37°C in an aerobic atmosphere with 5% CO₂ on Middlebrook 7H10 agar with OADC enrichment

⁵NR-49073 was deposited as *M. florentinum* and reported to be negative for growth in presence of 5% sodium chloride. Testing performed in triplicate by BEI Resources indicates a positive result

⁶Relatedness between bacterial strains has traditionally been determined using DDH. For additional information, refer to Auch, A.F., et al. "Digital DNA-DNA Hybridization for Microbial Species Delineation by Means of Genome-to-Genome Sequence Comparison." *Stand. Genomic Sci.* 2 (2010): 117-134. PubMed: 21304684.

Certificate of Analysis for NR-49073

⁷The whole genome of *M. florentinum*, strain FI-93171T (Contig Total length 6.2 megabase pairs) was sequenced using the Illumina® MiSeq® system and was assembled and analyzed with CLC Genomics Workbench Version 7.0.2.

⁸The required whole genome sequence for the type strain of this species is not available. dDDH testing rules out all species listed in Table 1, however, this does not rule out species for which the type strains whole genome sequences are not available.

⁹Purity of this lot was assessed for 14 days at 37°C in an aerobic atmosphere with 5% CO₂.

Figure 1: Colony Morphology**Table 1: Digital DNA-DNA hybridization (dDDH)**

Species	Strain	Accession #	GGD vs. NR-49073 (Deposited as: <i>M. florentinum</i>)
<i>M. abscessus</i> subsp. <i>abscessus</i>	Hauduroy 948 ^T	NC_010397.1	19.1
<i>M. abscessus</i> subsp. <i>bolletii</i>	BD ^T	AHAS00000000.1	19
<i>M. abscessus</i> subsp. <i>massiliense</i>	CCUG 48898 ^T	NZ_AP014547.1	19
<i>M. aromaticivorans</i>	JS19b1 ^T	JALN00000000.2	19.9
<i>M. aurum</i>	ATCC 23366 ^T	CVQQ01000001.1	20.2
<i>M. austroafricanum</i>	E9789-SA12441 ^T	HG964450.1	19.6
<i>M. avium</i> subsp. <i>avium</i>	ATCC 25291 ^T	ACFI00000000.1	24.2
<i>M. avium</i> subsp. <i>paratuberculosis</i>	ATCC 19698 ^T	AGAR00000000.1	24.7
<i>M. avium</i> subsp. <i>silvaticum</i>	6409 ^T	AYOC00000000.1	24.7
<i>M. bohemicum</i>	CIP 105808 ^T	CSTD01000001.1	23.4
<i>M. canariense</i>	502329 ^T	BCSY00000000.1	20.1
<i>M. celatum</i>	ATCC 51131 ^T	BBUN00000000.1	22.2
<i>M. chelonae</i>	CM 6388 ^T	CP010946.1	19.6
<i>M. chlorophenicolum</i>	PCP-I ^T	JYNL00000000.1	20
<i>M. chubuense</i>	48013 ^T	NC_018027.1	19.9
<i>M. colombiense</i>	10B ^T	AFVW00000000.2	24.1
<i>M. conceptionense</i>	D16 ^T	CTEF00000000.1	20.2
<i>M. cosmeticum</i>	LTA-388 ^T	CCBB000000000.1	20
<i>M. crocinum</i>	czh-42 ^T	BBHD00000000.1	21.4
<i>M. färncinogenes</i>	IEMVT 75 ^T	CCAY000000000.1	20
<i>M. fluoranthenivorans</i>	FA4 ^T	BBFT00000000.1	21.1
<i>M. fortuitum</i> subsp. <i>fortuitum</i>	ATCC 6841 ^T	CP014258.1	19.7
<i>M. fortuitum</i> subsp. <i>acetamidolyticum</i>	NCH E11620 ^T	BCSZ00000000.1	19.6
<i>M. gastri</i>	ATCC 15754 ^T	AZYN00000000.1	22.4
<i>M. genavense</i>	2289 ^T	JAGZ00000000.1	34.1
<i>M. haemophilum</i>	ATCC 29548 ^T	CP011883.2	22.6
<i>M. hawaiiicum</i>	3849 ^T	ARBU00000000.1	20.3
<i>M. hodleri</i>	EMI2 ^T	BBGO00000000.1	22.9
<i>M. intracellularare</i>	ATCC 13950 ^T	NC_016946.1	24.1

Certificate of Analysis for NR-49073

Species	Strain	Accession #	GGD vs. NR-49073 (Deposited as: <i>M. florentinum</i>)
<i>M. kansasii</i>	ATCC 12478 ^T	NC_022663.1	22.1
<i>M. kyorinense</i>	KUM 060204 ^T	BBKA00000000.1	22
<i>M. mageritense</i>	938 ^T	CCBF00000000.1	20
<i>M. neoaurum</i>	ATCC 25795 ^T	JMDW00000000.1	19.8
<i>M. neworleansense</i>	W6705 ^T	CWKH00000000.1	19.7
<i>M. novocastrense</i>	73 ^T	BCTA00000000.1	20.2
<i>M. obuense</i>	47001 ^T	JYNU00000000.1	19.6
<i>M. pallens</i>	czh-8 ^T	BBHE00000000.1	21.6
<i>M. parascrofulaceum</i>	HSC-68 ^T	ADNV00000000.1	24.3
<i>M. pseudoshottsii</i>	L15 ^T	BCND00000000.1	21.4
<i>M. pyrenivorans</i>	17A3 ^T	BBHB00000000.1	22.1
<i>M. rufum</i>	JS14 ^T	JROA00000000.1	19.9
<i>M. rutilum</i>	czh-117 ^T	BBHF00000000.1	23.3
<i>M. septicum</i>	W4964 ^T	CBMO00000000.1	19.9
<i>M. setense</i>	ABO-M06 ^T	JTJW00000000.1	19.6
<i>M. simiae</i>	ATCC 25275 ^T	CBMJ00000000.2	26.5
<i>M. smegmatis</i>	ATCC 19420 ^T	LN831039.1	20.1
<i>M. thermoresistibile</i>	ATCC 19527 ^T	BCTB00000000.1	20
<i>M. triplex</i>	90-1019 ^T	CCAU00000000.1	36.7
<i>M. tuberculosis</i>	H37Rv ^T	NC_000962.3	22.1
<i>M. vaccae</i>	ATCC 15483 ^T	BCRS00000000.1	20.1
<i>M. vanbaalenii</i>	PYR-1 ^T	NC_008726.1	20.1
<i>M. vulneris</i>	NLA000700772 ^T	CCBG00000000.1	19.9
<i>M. yongonense</i>	05-1390 ^T	NC_021715.1	24.1
<i>Nocardia asteroides</i>	NBRC 15531 ^T	BAFO00000000.2	19.6

Date: 23 JUN 2017

Signature:



BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

