

Certificate of Analysis for NR-49368

Mycobacterium tuberculosis, Strain XTB13-092

Catalog No. NR-49368

Product Description: *Mycobacterium tuberculosis* (*M. tuberculosis*), strain XTB13-092 was isolated in 2011 from the sputum of a patient with tuberculosis in the Republic of Belarus. Strain XTB13-092 was deposited as resistant to isoniazid, rifampin and streptomycin.

Lot¹: 64064223 Manufacturing Date: 11MAY2016

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis ²		
Cellular morphology	Gram-positive rods	Gram-positive rods
Colony morphology ³	Report results	Irregular, slight peaked, undulate, rough and cream (Figure 1)
Growth rate	≥ 7 days	21 days
Growth at 26°C	Negative	Negative
Growth at 37°C	Positive	Positive
Acid-fast stain	Positive (red colonies)	Positive (red colonies)
Pigmentation in the dark (Scotochromogen)	Negative (no pigment)	Negative (no pigment)
Photoinduction for 1 hour (Photochromogen)	Negative (no pigment)	Negative (no pigment)
Nonchromogen (no pigment)	Positive (no pigment)	Positive (no pigment)
Biochemical tests	, , ,	, , , ,
Niacin production ⁴	Positive	Positive
Nitrate reduction	Positive	Positive
Pyrazinamidase	Positive	Positive
Antibiotic Susceptibility Profile Sensititre™ System ^{5,6}		
Amikacin	Report results	0.25 μg/mL
Cycloserine	Report results	8 μg/mL
Ethambutol	Report results	1 μg/mL ⁷
Ethionamide	Report results	5 μg/mL ⁷
Isoniazid	Report results	4 μg/mL
Kanamycin	Report results	2.5 μg/mL
Moxifloxacin	Report results	0.25 µg/mL
Ofloxacin	Report results	1 μg/mL
Para-aminosalicylic acid	Report results	2 μg/mL ⁷
Rifabutin	Report results	16 µg/mL ⁷
Rifampin	Report results	16 μg/mL_
Streptomycin	Report results	32 μg/mL ⁷
Genotypic Analysis		
Sequencing of Heat Shock Protein 65 gene (~ 430 base pairs)	≥ 99% sequence identity to <i>M. tuberculosis</i> , strain XTB13-092 (GenBank: JLMB01000001.1)	100% sequence identity to M. tuberculosis, strain XTB13-092 (GenBank: JLMB01000001.1)8
Purity (post-freeze)		
Middlebrook 7H10 agar with OADC enrichment ⁹	Growth consistent with expected	Growth consistent with expected
Č	colony morphology	colony morphology
Tryptic Soy agar ¹⁰	Report results	Growth consistent with expected colony morphology
Viability (post-freeze) ³	Growth	Growth

¹NR-49368 was produced by inoculation of the deposited material into Middlebrook 7H9 broth with ADC enrichment. Broth inoculum was added to Middlebrook 7H10 agar with OADC enrichment kolles, which were grown for 62 days at 37°C in an aerobic atmosphere with 5% CO₂ to produce this lot.

BEI Resources

www.beiresources.org

E-mail: contact@beiresources.org

Tel: 800-359-7370 Fax: 703-365-2898



Certificate of Analysis for NR-49368

mycobacteria and Lévy-Frébault, V. V. and F. Portaels. "Proposed Minimal Standards for the Genus Mycobacterium and for Description of New Slowly Growing Mycobacterium Species." Int. J. Syst. Bacteriol. 42 (1992): 315-323. PubMed: 1581193.

³21 days at 37°C in an aerobic atmosphere with 5% CO₂ on Middlebrook 7H10 agar with OADC enrichment

⁴All mycobacteria produce niacin but only *M. tuberculosis* accumulates it, resulting in a positive test for *M. tuberculosis*.

⁸Also consistent with *M. africanum*, *M. bovis*, *M. canettii*, *M. caprae* and *M. microti*

¹⁰Purity of this lot was assessed for 21 days at 37°C in an aerobic atmosphere with 5% CO₂.





Date: 26 OCT 2017 Signature:

BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

BEI Resources

www.beiresources.org

E-mail: contact@beiresources.org

Tel: 800-359-7370 Fax: 703-365-2898

⁵Sensititre[™] System *Mycobacterium tuberculosis* MIC Plate, Thermo Scientific[™], catalog number MYCOTB ⁶Minimum Inhibitory Concentration (MIC); No Clinical & Laboratory Standards Institute (CLSI) interpretations of the Sensititre[™] System data for M. tuberculosis are currently available.

⁷For streptomycin, ethionamide, para-aminosalicylic acid, rifabutin and ethambutol, the endpoint for these drugs is determined by the well with approximately 80% inhibition of growth compared to the positive control well with no drug.

⁹Purity of this lot was assessed for 51 days at 37°C in an aerobic atmosphere with 5% CO₂.