

***Klebsiella pneumoniae*, Strain BIDMC 12A**

Catalog No. NR-41928

Product Description: *Klebsiella pneumoniae* (*K. pneumoniae*), strain BIDMC 12A was isolated in 2012 from human sputum in Boston, Massachusetts, USA.¹ *K. pneumoniae*, strain BIDMC 12A was deposited as a carbapenem-resistant strain and is part of a Carbapenem-Resistant Enterobacteriaceae (CRE) Sequencing Project at the Broad Institute. Strain BIDMC 12A was deposited as resistant to ampicillin/sulbactam, cefepime, meropenem and trimethoprim/sulfamethoxazole.

Lot¹: 70007985

Manufacturing Date: 11AUG2017

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis Cellular morphology Colony morphology ² Motility (wet mount) VITEK® 2 Compact (GN card)	Gram-negative rods Report results Report results <i>K. pneumoniae</i> (≥ 89%)	Gram-negative rods Circular, convex, entire, mucoid and cream (Figure 1) Non-motile <i>K. pneumoniae</i> (99%) ³
Antibiotic Susceptibility Profile VITEK® (AST-GN83) ⁴ Ampicillin Amoxicillin/clavulanic acid Ampicillin/sulbactam Piperacillin/tazobactam Cefazolin Cefuroxime Cefuroxime axetil Cefoxitin Cefotaxime Ceftazidime Ceftriaxone Cefepime Aztreonam Meropenem Amikacin Gentamicin Ciprofloxacin Nitrofurantoin Trimethoprim/sulfamethoxazole Etest® antibiotic test strips ⁵ Ceftriaxone ⁶ Tobramycin ⁶	Report results Report results Resistant Report results Report results Report results Report results Report results Report results Report results Report results Resistant Report results Resistant Report results Report results Report results Report results Resistant Report results Report results Report results Resistant Report results Report results	Resistant (≥ 32 µg/mL) Resistant (≥ 32 µg/mL) Resistant (≥ 32 µg/mL) Resistant (≥ 128 µg/mL) Resistant (≥ 64 µg/mL) Resistant (≥ 64 µg/mL) Resistant (≥ 64 µg/mL) Resistant (≥ 32 µg/mL) Resistant (= 8 µg/mL) Resistant (≥ 64 µg/mL) Resistant (≥ 64 µg/mL) Resistant (= 8 µg/mL) Resistant (≥ 64 µg/mL) Resistant (≥ 16 µg/mL) Resistant (≥ 64 µg/mL) Sensitive (= 4 µg/mL) Resistant (≥ 4 µg/mL) Resistant (= 256 µg/mL) Resistant (≥ 320 µg/mL) Resistant (= 32 µg/mL) Intermediate (= 8 µg/mL)
Genotypic Analysis Sequencing of 16S ribosomal RNA gene (~ 710 base pairs)	≥ 99% sequence identity to <i>K. pneumoniae</i> , strain BIDMC 12A (GenBank: JANE01000004.1)	99.6% sequence identity to <i>K. pneumoniae</i> , strain BIDMC 12A (GenBank: JANE01000004.1) ⁷
Purity (post-freeze)⁸	Growth consistent with expected colony morphology	Growth consistent with expected colony morphology
Viability (post-freeze)²	Growth	Growth

¹NR-41928 was produced by inoculation of the deposited material into Tryptic Soy broth and grown for 1 day at 37°C in an aerobic atmosphere. Broth inoculum was added to Tryptic Soy agar kolles, which were grown for 1 day at 37°C in an aerobic atmosphere to produce this lot.

²1 day at 37°C in an aerobic atmosphere on Tryptic Soy agar

³Percent probabilities above 89% indicate a close match to the typical biochemical pattern for the given organism, with a percent probability of 99% being a perfect match between the test reaction pattern and the unique biochemical pattern of the given organism or organism group. For additional information, please refer to O'Hara, C. M. and J. M. Miller. "Evaluation of the VITEK 2 ID-GNB Assay for Identification of Members of the Family Enterobacteriaceae and Other Nonenteric Gram-Negative Bacilli and Comparison with the VITEK GNI+ Card." *J. Clin. Microbiol.* 41 (2003): 2096-2101. PubMed: 12734254.

⁴Minimum Inhibitory Concentration (MIC); MIC interpretation was determined using VITEK[®] 2 software version 07.01 combined with the bioMérieux Advanced Expert System[™] (AES) software using the interpretation standard CLSI M100-S22 (2012) and the interpretation guideline "Natural Resistance." For more information, please refer to Sanders, C. C., et al. "Potential Impact of the VITEK 2 System and the Advanced Expert System on the Clinical Laboratory of a University-Based Hospital." *J. Clin. Microbiol.* 39 (2001): 2379-2385. PubMed: 11427542.


⁵1 day at 37°C in an aerobic atmosphere on Mueller Hinton agar

⁶MIC Interpretation Guideline: CLSI M100-S22 (2012)

⁷Also consistent with other organisms

⁸Purity of this lot was assessed for 7 days at 37°C in an aerobic atmosphere with 5% CO₂ on Tryptic Soy agar.

Figure 1: Colony Morphology



Date: 05 JAN 2018

Signature:

BEI Resources Authentication

ATCC[®], on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC[®]'s knowledge.

ATCC[®] is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

