SUPPORTING INFECTIOUS DISEASE RESEARCH

Mycobacterium tuberculosis, Strain 98-3182

Catalog No. NR-30904

This reagent is the tangible property of the U.S. Government.

Product Description: *Mycobacterium tuberculosis (M. tuberculosis)*, strain 98-3182 was isolated between 1995 and 2000 from human sputum from an HIV-negative patient infected with pulmonary tuberculosis in North America.

Lot¹: 70004570

Manufacturing Date: 02JUN2017

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis ²		
Cellular morphology	Gram-positive rods	Gram-positive rods
Colony morphology ³	Report results	Irregular, slight peaked, undulate, rough and cream (Figure 1)
Growth rate	≥ 7 days	21 days
Growth at 26°C	Negative	Negative
Growth at 37°C	Positive	Positive
Acid-fast stain	Positive (red colonies)	Positive (red colonies)
Pigmentation in the dark (Scotochromogen)	Negative (no pigment)	Negative (no pigment)
Photoinduction for 1 hour (Photochromogen)	Negative (no pigment)	Negative (no pigment)
Nonchromogen (no pigment) Biochemical tests	Positive (no pigment)	Positive (no pigment)
Niacin production ⁴	Positive	Positive
Nitrate reduction	Positive	Positive
Pyrazinamidase	Positive	Positive
Antibiotic Susceptibility Profile Sensititre™ System ^{5,6}		
Amikacin	Report results	0.25 µg/mL
Cycloserine	Report results	0.25 μg/mL 16 μg/mL
Ethambutol	Report results	$1 \mu\text{g/mL}^7$
Ethionamide	Report results	$1.2 \mu\text{g/mL}^7$
Isoniazid	Report results	0.06 μg/mL
Kanamycin	Report results	2.5 μg/mL
Moxifloxacin	Report results	0.25 μg/mL
Ofloxacin	Report results	1 μg/mL
Para-aminosalicylic acid	Report results	≤ 0.5 μg/mL ⁷
Rifabutin	Report results	$\leq 0.12 \ \mu g/mL^7$
Rifampin	Report results	≤ 0.12 µg/mL
Streptomycin	Report results	$0.5 \ \mu g/mL^7$
Genotypic Analysis		r or
Sequencing of Heat Shock Protein 65 gene	≥ 99% sequence identity to	100% sequence identity to
(~ 410 base pairs)	<i>M. tuberculosis</i> type strain (GenBank: AL123456)	<i>M. tuberculosi</i> s type strain (GenBank: AL123456) ⁸
Purity (post-freeze)		
Middlebrook 7H10 agar with OADC enrichment ⁹	Growth consistent with expected colony morphology	Growth consistent with expected colony morphology
Tryptic Soy agar ¹⁰	Report results	Growth consistent with expected colony morphology
Viability (post-freeze) ³	Growth	Growth

¹NR-30904 was produced by inoculation of the deposited material into Middlebrook 7H9 broth with ADC enrichment. Broth inoculum was added to Middlebrook 7H10 agar with OADC enrichment kolles, which were grown for 21 days at 37°C in an aerobic atmosphere with 5% CO₂ to produce this lot.

BEI Resources www.beiresources.org E-mail: <u>contact@beiresources.org</u> Tel: 800-359-7370 Fax: 703-365-2898 dei resources

Certificate of Analysis for NR-30904

SUPPORTING INFECTIOUS DISEASE RESEARCH

²Information on Mycobacterium testing is available from Ribón, W. "Biochemical Isolation and Identification of Mycobacteria." <u>Biochemical Testing</u>. (2012) Jose C. Jimenez-Lopez (Ed.), InTech, <u>http://www.intechopen.com/books/biochemical-testing/biochemical-isolation-and-identification-of-mycobacteria</u> and Lévy-Frébault, V. V. and F. Portaels. "Proposed Minimal Standards for the Genus *Mycobacterium* and for Description of New Slowly Growing *Mycobacterium* Species." <u>Int. J. Syst. Bacteriol.</u> 42 (1992): 315-323. PubMed: 1581193.

³21 days at 37°C in an aerobic atmosphere with 5% CO₂ on Middlebrook 7H10 agar with OADC enrichment

⁴All mycobacteria produce niacin but only *M. tuberculosis* accumulates it, resulting in a positive test for *M. tuberculosis*.

⁵SensititreTM System *Mycobacterium tuberculosis* MIC Plate, Thermo ScientificTM, catalog number MYCOTB

⁶Minimum Inhibitory Concentration (MIC); No Clinical & Laboratory Standards Institute (CLSI) interpretations of the Sensititre™ System data for *M. tuberculosis* are currently available.

⁷For streptomycin, ethionamide, para-aminosalicylic acid, rifabutin and ethambutol, the endpoint for these drugs is determined by the well with approximately 80% inhibition of growth compared to the positive control well with no drug.

⁸Also consistent with *M. africanum*, *M. bovis*, *M. canettii*, *M. caprae* and *M. microti*

⁹Purity of this lot was assessed for 39 days at 37°C in an aerobic atmosphere with 5% CO₂.

¹⁰Purity of this lot was assessed for 21 days at 37°C in an aerobic atmosphere with 5% CO₂.

Figure 1: Colony Morphology

/Heather Couch/ Heather Couch

<u>27 JUN 2018</u>

Program Manager or designee, ATCC Federal Solutions

ATCC[®], on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC[®]'s knowledge.

ATCC[®] is a trademark of the American Type Culture Collection. You are authorized to use this product for research use only. It is not intended for human use.

