

Simian Virus 40, Baylor (SVB2E-WT)

Catalog No. NR-51200

Product Description: SV40, Baylor (SVB2E-WT) was isolated in 1961 from a type 2 Sabin oral polio vaccine prepared in 1956.

Passage History: AGMK7CV7TC3/CV1 (Prior to deposit at BEI Resources/BEI Resources); AGMK = African Green monkey kidney cells; CV = CV-1 cells¹; TC = TC-7 cells

Lot²: 70016132

Manufacturing Date: 16JUL2018

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity in CV-1 cells	Cell rounding, vacuolization and detachment	Cell rounding, vacuolization and detachment
Sequencing of Species-Specific Region (~ 960 nucleotides)	≥ 98% identity with SV40, B-2 (GenBank: AF155358.1)	100% identity with SV40, B-2 (GenBank: AF155358.1)
Titer by TCID₅₀ Assay^{3,4} in CV-1 cells¹ by Cytopathic Effect	Report results	2.8 × 10 ⁷ TCID ₅₀ per mL
Amplification of SV40 Sequence by PCR	~ 1080 base pair amplicon	~ 1080 base pair amplicon
Sterility (21-day incubation) Harpo's HTYE broth ⁵ , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Blood agar, 37°C, aerobic Blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO ₂	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA detection by PCR of extracted Test Article nucleic acid	None detected None detected	None detected None detected

¹CV-1: ATCC® CCL-70™

²Grown in Eagle's Minimum Essential Medium containing Earle's Balanced Salt Solution, non-essential amino acids, 2 mM L-glutamine, 1 mM sodium pyruvate and 1.5 g/L of sodium bicarbonate (ATCC® 30-2003) supplemented with 2% fetal bovine serum (ATCC® 30-2020) for 11 days at 37°C with 5% CO₂

³The Tissue Culture Infectious Dose 50% (TCID₅₀) endpoint is the 50% infectious endpoint in cell culture. The TCID₅₀ is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID₅₀ provides a measure of the titer (or infectivity) of a virus preparation.

⁴Assay plates were incubated 9 days at 37°C and 5% CO₂

⁵Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798

/Heather Couch/

Heather Couch

05 DEC 2018

Program Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

