

***Leishmania mexicana*, Strain MNYC/BZ/62/M379 (+*luc*)**

**Catalog No. NR-51210**

**Product Description:** *Leishmania mexicana* (*L. mexicana*), strain MNYC/BZ/62/M379 (+*luc*) is a transgenic clone derived from the parent MNYC/BZ/62/M379 strain. Strain MNYC/BZ/62/M379, also referred to as M 379 or MNYZ/BZ/62/M379, was originally isolated in 1962 from a *Nyctomys sumichrasti* (vesper rat) in Cayo District, Belize (formerly British Honduras).

**Lot<sup>1</sup>: 70014487**

**Manufacturing Date: 05APR2018**

TEST	SPECIFICATIONS	RESULTS
<b>Cellular Morphology<sup>2</sup></b>	Report results	Elongated, motile, refractile
<b>Genotyping<sup>3</sup></b> Sequencing of internal transcribed spacer (ITS) 1, 5.8S ribosomal RNA gene, ITS 2 (~ 1010 base pairs)	≥ 99% sequence identity to <i>L. mexicana</i> , strain MNYC/BZ/62/M379 (GenBank: AF466383.1)	99.0% sequence identity to <i>L. mexicana</i> , strain MNYC/BZ/62/M379 (GenBank: AF466383.1)
<b>Functional Activity by PCR Amplification<sup>3</sup></b> ITS 1, 5.8S rRNA gene, ITS 2	~ 1200 base pair amplicon	~ 1200 base pair amplicon
<b>Functional Activity of Luciferase Gene<sup>3,4</sup></b>	Positive	Positive
<b>Viable Cell Count by Hemacytometry<sup>3</sup></b>	> 10 <sup>6</sup> cells per mL	5.4 × 10 <sup>8</sup> cells per mL
<b>Viability (post-freeze)<sup>2,5</sup></b>	Growth	Growth
<b>Sterility (21-day incubation)<sup>2</sup></b> Harpo's HTYE broth <sup>6</sup> , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic DMEM with 10% FBS, 37°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth

<sup>1</sup>NR-51210 was produced by cultivation of the deposited material in Modified M199 medium supplemented with 10% heat-inactivated fetal bovine serum and 10 µg/mL hemin for 3 days at 25°C in an aerobic atmosphere to produce this lot.

<sup>2</sup>Testing completed on vial, post-freeze material.

<sup>3</sup>Testing completed on bulk material prior to vialing and freezing.

<sup>4</sup>Luciferase activity was determined using the Promega *Renilla* Luciferase Assay System (Catalog No. E2810). Parasites were lysed and incubated with luciferase assay reagent. Light was measured using a spectrophotometer with a wavelength of 450 nm.

<sup>5</sup>Viable cells were observed after 2 days in Modified M199 medium supplemented with 10% heat-inactivated fetal bovine serum and 10 µg/mL hemin at 25°C in an aerobic atmosphere.

<sup>6</sup>Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

/Heather Couch/  
Heather Couch

27 FEB 2019

Program Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.  
You are authorized to use this product for research use only. It is not intended for human use.

