

***Ehrlichia chaffeensis*, Strain Jax**

Catalog No. NR-46444

This reagent is the property of the U.S. Government

Product Description: Cell lysate and supernatant from *Canis familiaris* macrophage-monocyte cells¹ infected with *Ehrlichia chaffeensis* (*E. chaffeensis*), strain Jax, containing 20% fetal bovine serum and 10% DMSO (final concentrations).

Lot²: 62921182

Manufacturing Date: 21FEB2015

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity in DH82 Cells ¹	Report results	Positive detection of morulae formation (Diff-Quick stain)
Identification by Sequencing of Species-Specific Region (826 nucleotides)	Consistent with <i>E. chaffeensis</i> , strain Jax	100% identity with <i>E. chaffeensis</i> , strain Jax (GenBank: CP007475)
Titer by TCID ₅₀ Assay ^{3,4} in DH82 Cells ¹ by IFA ⁵	Report results	1.6 × 10 ⁶ TCID ₅₀ per mL
Sterility (21-day incubation) Harpo's HTYE broth ⁶ , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Brucella agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO ₂	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA detection by PCR of extracted Test Article nucleic acid	None detected None detected	None detected None detected

¹DH82 cells: ATCC® CRL-10389™

²Grown in Dulbecco's Modified Eagle's Medium containing 4 mM L-glutamine, 4500 mg per L glucose, 1 mM sodium pyruvate, and 1500 mg per L sodium bicarbonate (ATCC® 30-2002), supplemented with 5% fetal bovine serum (ATCC® 30-2020) for 11 days at 37°C with 5% CO₂

³The Tissue Culture Infectious Dose 50% (TCID₅₀) endpoint is the 50% infectious endpoint in cell culture. The TCID₅₀ is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID₅₀ provides a measure of the titer (or infectivity) of a virus preparation.


⁴10 days at 37°C and 5% CO₂

⁵Using *Ehrlichia chaffeensis* IFA IgG reagent kit (Fuller Laboratories ECHG-120)

⁶Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

Date: 06 APR 2016

Signature:



BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

