

**Enterovirus 71 (EV-71), Tainan/4643/1998**

**Catalog No. NR-471**

**Product Description:**

Cell lysate and supernatant from human rhabdomyosarcoma (RD) cells infected with EV-71, Tainan/4643/1998.<sup>1</sup>

**Lot: 7746372<sup>2</sup>**

**Manufacturing Date: 25JAN2007**

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity in RD Cells <sup>1</sup>	Report results	Cell rounding and detachment
Identification by Indirect Fluorescent Antibody Assay Monoclonal antibody to EV-71 <sup>3</sup>	Fluorescence observed	Fluorescence observed
Sequencing of a Strain-Specific Region (~ 530 nucleotides)	Report results	Identical to GenBank AF304458
RT-PCR Amplification of an Enterovirus-Specific Region	~ 680 base pair amplicon	~ 680 base pair amplicon
Titer by TCID <sub>50</sub> Assay in RD Cells <sup>1,4,5</sup>	Report results	1.6 × 10 <sup>8</sup> TCID <sub>50</sub> /mL
<b>Sterility (21-day incubation)</b> Harpo's HTYE broth, 37°C and 26°C, aerobic <sup>6</sup> Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO <sub>2</sub>	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
<b>Mycoplasma Contamination</b> Agar and broth culture (14-day incubation at 37°C) DNA Detection by PCR of Test Article nucleic acid	None detected None detected	None detected None detected

<sup>1</sup>RD cells: ATCC® CCL-136™ (lot 4083062)

<sup>2</sup>EV-71, Tainan/4643/1998 was deposited by National Cheng Kung University, Tainan, Taiwan. NR-471 was grown from the deposited virus seed in Minimum Essential Medium containing Earle's salts and non-essential amino acids (Invitrogen™ 10370) supplemented with 2% irradiated fetal bovine serum (Cambrex® 14-471F), 2 mM L-glutamine (Invitrogen™ 25030), and 1 mM sodium pyruvate (Invitrogen™ 11360) for 3 days at 37°C and 5% CO<sub>2</sub>.

<sup>3</sup>Millipore MAB979

<sup>4</sup>The Tissue Culture Infectious Dose 50% (TCID<sub>50</sub>) endpoint is the 50% infectious endpoint in cell culture. The TCID<sub>50</sub> is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD<sub>50</sub>) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID<sub>50</sub> provides a measure of the titer (or infectivity) of a virus preparation.

<sup>5</sup>6 days at 37°C and 5% CO<sub>2</sub>.

<sup>6</sup>Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

/Heather Couch/

Heather Couch

24 FEB 2020

Program Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

