

Spike Glycoprotein (Stabilized) from SARS-Related Coronavirus 2, Wuhan-Hu-1 with C-Terminal Histidine Tag, Recombinant from HEK293F Cells

Catalog No. NR-52397

This reagent is the tangible property of the U.S. Government.

Product Description:

A recombinant form of the spike (S) glycoprotein from severe acute respiratory syndrome-related coronavirus 2 (SARS-CoV-2), Wuhan-Hu-1 (GenPept: [QJE37812](#)) was produced by transfection of purified plasmid (BEI Resources NR-52394 lot 70034724) in human embryonic kidney HEK293F (Expi293F™; Gibco™ A14527) cells, purified by nickel affinity (HiTrap™ HP) chromatography and vialled in phosphate-buffered saline (PBS), pH 7.4. NR-52397 lacks the signal sequence and contains 1196 residues (ectodomain) of the SARS-CoV-2 spike glycoprotein; the recombinant protein was modified to remove the polybasic S1/S2 cleavage site (RRAR to A; residues 682 to 685), stabilized with a pair of mutations (K986P and V987P, wild type numbering; GenPept: [YP_009724390](#)) and includes a thrombin cleavage site, T4 foldon trimerization domain and C-terminal hexa-histidine tag.

Lot: 70035631


Manufacturing Date: 26MAY2020

TEST	SPECIFICATIONS	RESULTS
Appearance	Clear and colorless	Clear and colorless
SDS-PAGE Analysis	Protein band of interest represents > 90% of total staining intensity	Protein band of ~ 195 kDa represents > 95% of total staining intensity (Figure 1) ¹
Concentration by Bicinchoninic Acid Assay Bovine Serum Albumin (standard)	Report results	0.38 mg per mL
Final Product Amount per vial Volume per vial	Report results Report results	45 µg 118 µL
Functional Activity by Western Blot Analysis Monoclonal anti-histidine tag	Reactive	Reactive (Figure 2) ²
Sterility	0.22 µm sterile-filtered	0.22 µm sterile-filtered

¹The recombinant protein migrated to a slightly larger size than was expected, likely caused by glycosylation common in recombinant spike proteins derived from coronaviruses. For more information, please see Chakraborti, S., et al. "The SARS Coronavirus S Glycoprotein Receptor Binding Domain: Fine Mapping and Functional Characterization." *Virology*, 2 (2005): 73. PubMed: 16122388.


²Using a 1:1000 dilution of mouse monoclonal anti-histidine tag [Santa Cruz Biotechnology, Inc., His-probe (H-3): sc-8036 HRP] as primary antibody.

Figure 1: SDS-PAGE Analysis



Lane 1: Precision Plus Protein™ Standard (8 µL)
Lane 2: NR-52397 (2 µg)
Lane 3: Bovine serum albumin (6 µg)

Figure 2: Anti-Histidine Western Blot Analysis



Lane 1: MagicMark™ XP Protein Standard (8 µL)
Lane 2: NR-52397 (0.5 µg)
Lane 3: Bovine serum albumin (1.5 µg)

/Heather Couch/
Heather Couch

11 JUN 2020

Program Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected by the contributor to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

