

**Spike Glycoprotein Receptor Binding Domain (RBD) from SARS-Related Coronavirus 2, Wuhan-Hu-1 with C-Terminal Histidine Tag, Recombinant from Baculovirus**

**Catalog No. NR-52307**

This reagent is the tangible property of the U.S. Government.

**Product Description:**

A recombinant form of the spike glycoprotein receptor binding domain (RBD) from severe acute respiratory syndrome-related coronavirus 2 (SARS-CoV-2), Wuhan-Hu-1 (GenPept: [QHD43416](#)) was produced in Sf9 insect cells using a baculovirus expression system and purified by nickel affinity and ion exchange chromatography. NR-52307 contains 223 residues of the SARS-CoV-2 spike glycoprotein RBD and features a C-terminal hexa-histidine tag.

**Lot: 70034256**


**Manufacturing Date: 13MAY2020**

| TEST   | SPECIFICATIONS  | RESULTS   |
|--|---|---|
| <b>Appearance</b>  | Clear and colorless   | Clear and colorless   |
| <b>SDS-PAGE Analysis (Coomassie Blue)</b>  | Protein band of interest represents > 90% of total staining intensity | Protein band of ~ 31 kDa represents > 90% of total staining intensity (Figure 1) <sup>1</sup> |
| <b>Concentration by Bicinchoninic Acid Assay</b><br>Bovine Serum Albumin (standard)  | Report results  | 0.103 mg per mL   |
| <b>Final Product</b><br>Amount per vial<br>Volume per vial                           | Report results<br>Report results                                      | 29 µg<br>280 µL   |
| <b>Functional Activity by Western Blot Analysis</b><br>Monoclonal anti-histidine tag | Reactive  | Reactive (Figure 2) <sup>2</sup>  |
| <b>Sterility</b>   | 0.2 µm sterile-filtered   | 0.2 µm sterile-filtered   |

<sup>1</sup>The recombinant protein migrated to a slightly larger size than was expected, likely caused by glycosylation common in recombinant spike proteins derived from coronaviruses. For more information, please see Chakraborti, S., et al. "The SARS Coronavirus S Glycoprotein Receptor Binding Domain: Fine Mapping and Functional Characterization." *Virology*, 2 (2005): 73. PubMed: 16122388.


<sup>2</sup>Using a 1:1000 dilution of mouse monoclonal anti-histidine tag (Clontech 631212) as primary antibody and a 1:1000 dilution of HRP-conjugated goat anti-mouse IgG (R&D Systems HAF007) as secondary antibody.

**Figure 1: SDS-PAGE Analysis**



Lane 1: Precision Plus Protein™ Standard (6 µL)  
Lane 2: NR-52307 (1 µg)  
Lane 3: Bovine serum albumin (1 µg)

**Figure 2: Anti-Histidine Western Blot Analysis**



Lane 1: MagicMark™ XP Protein Standard (5 µL)  
Lane 2: NR-52307 (0.1 µg)  
Lane 3: Bovine serum albumin (0.1 µg)

/Heather Couch/

Heather Couch

28 JUL 2020

Program Manager or designee, ATCC Federal Solutions

ATCC<sup>®</sup>, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC<sup>®</sup>'s knowledge.

ATCC<sup>®</sup> is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

