

**Enterovirus Species D Type 68, USA/2009-23228**

**Catalog No. NR-51993**

**Product Description:**

Enterovirus species D type 68 (EV-D68), USA/2009-23228 was isolated in 2009 from a respiratory sample from a human subject in Indiana, USA. The subject was not suffering from acute flaccid myelitis (AFM). NR-51993 lot 70040493 was produced by infecting rhabdomyosarcoma cells (RD; ATCC® CCL-136™) and incubating in Eagle's Minimum Essential Medium (ATCC® 30-2003™) supplemented with 2% fetal bovine serum (ATCC® 30-2020™) for 3 days at 33°C with 5% CO<sub>2</sub>.

**Passage History:**

RD(4)/RD(2) (Prior to deposit at BEI Resources/BEI Resources); RD = rhabdomyosarcoma cells

**Lot: 70040493**

**Manufacturing Date: 10JAN2021**

TEST	SPECIFICATIONS	RESULTS
<b>Identification by Infectivity in RD Cells</b>	Cell rounding and detachment	Cell rounding and detachment
<b>Next-Generation Sequencing (NGS) of Complete Genome Using Illumina® iSeq™ 100 Platform</b> (~ 7300 nucleotides)	≥ 98% identity with EV-D68, USA/2009-23228 (GenBank: MN240504.1)	99.9% identity with EV-D68, USA/2009-23228 (GenBank: MN240504.1)
<b>Titer by TCID<sub>50</sub> Assay in RD Cells by Cytopathic Effect<sup>1</sup></b> (4 days at 33°C with 5% CO <sub>2</sub> )	Report results	1.6 × 10 <sup>7</sup> TCID <sub>50</sub> per mL
<b>Sterility (21-day incubation)</b> Harpo's HTYE broth, 37°C and 26°C, aerobic <sup>2</sup> Trypticase Soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C, aerobic	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
<b>Mycoplasma Contamination</b> Agar and broth culture (14-day incubation at 37°C) DNA detection by PCR of extracted Test Article nucleic acid	None detected None detected	None detected None detected

<sup>1</sup>The Tissue Culture Infectious Dose 50% (TCID<sub>50</sub>) endpoint is the 50% infectious endpoint in cell culture. The TCID<sub>50</sub> is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD<sub>50</sub>) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID<sub>50</sub> provides a measure of the titer (or infectivity) of a virus preparation.

<sup>2</sup>Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

/Heather Couch/  
Heather Couch

16 MAR 2021

Program Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

