

Antimicrobial Resistance Panel 8: *Pseudomonas aeruginosa* LpxC Inhibitor Resistant Mutants

Catalog No. NR-55647

Product Description:

NR-55647 consists of a 16-member panel of *Pseudomonas aeruginosa* (*P. aeruginosa*) strains containing mutations in the genes involved in the LpxC pathway. These strains were generated by selection on a growth medium containing an LpxC inhibitor or by introducing targeted mutations in the gene of interest.

The kit components were produced by inoculation of the deposited material into Tryptic Soy broth and grown for 1 day at 37°C in an aerobic atmosphere. Broth inoculums were added to Tryptic Soy agar kolles, which were grown for 1 day at 37°C in an aerobic atmosphere to produce the individual lots. Quality control testing was completed under propagation conditions unless otherwise noted.

BEI Resources is committed to ensuring digital accessibility for people with disabilities. This Certificate of Analysis contains complex tables and may not be fully accessible. Please let us know if you encounter accessibility barriers and a fully accessible document will be provided: E-mail: Contact@BEIResources.org. We try to respond to feedback within 24 hours.

Table 1: Kit Components

COMPONENT NUMBER	STRAIN	LOT NUMBER	MANUFACTURING DATE
NR-51885	<i>P. aeruginosa</i> , NB52019-CDA0033	70046530	18AUG2021
NR-51886	<i>P. aeruginosa</i> , NB52019-LpxCG208S	70046531	12AUG2021
NR-51887	<i>P. aeruginosa</i> , NB52019-CDJ0037	70048189	21OCT2021
NR-51888	<i>P. aeruginosa</i> , NB52217-P2	70046527	13AUG2021
NR-51889	<i>P. aeruginosa</i> , NB52217-P6	70046528	18AUG2021
NR-51890	<i>P. aeruginosa</i> , NB52217-P13	70046526	13AUG2021
NR-51891	<i>P. aeruginosa</i> , NB52217-PA4465 _{N193T}	70046529	13AUG2021
NR-51892	<i>P. aeruginosa</i> , NB52200-P6a (NC)	70046524	12AUG2021
NR-51893	<i>P. aeruginosa</i> , NB52200-P6b (SC)	70046525	13AUG2021
NR-51894	<i>P. aeruginosa</i> , NB52200-P13a (NC)	70046522	12AUG2021
NR-51895	<i>P. aeruginosa</i> , NB52200-P13b (SC)	70046523	12AUG2021
NR-51896	<i>P. aeruginosa</i> , NB52042-CDJ0042	70046532	13AUG2021
NR-51898	<i>P. aeruginosa</i> , NB52019-CDR0026	70046513	12AUG2021
NR-51899	<i>P. aeruginosa</i> , NB52019-CDR0061	70046514	18AUG2021
NR-51900	<i>P. aeruginosa</i> , NB52019-CDJ0011	70046516	13AUG2021
NR-51902	<i>P. aeruginosa</i> , NB52203-CDB0011	70048425	03NOV2021

Table 2: *Pseudomonas aeruginosa*, Strain NB52019-CDA0033 (NR-51885)

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis Cellular morphology Colony morphology Motility (wet mount) VITEK® MS (MALDI-TOF)	Gram-negative rods Report results Report results <i>P. aeruginosa</i>	Gram-negative rods Circular, convex, entire, smooth and cream Motile <i>P. aeruginosa</i> (99.9%)
Antibiotic Susceptibility Profile Etest® antibiotic test strips 1 day at 35°C in an aerobic atmosphere on Mueller Hinton agar Chloramphenicol Ofloxacin	Report results Report results	≥ 256 µg/mL ≥ 32 µg/mL
Genotypic Analysis Confirmation of mutations Digital DNA-DNA hybridization (dDDH) ¹	Mutations confirmed ≥ 70% for species identification	Pending <i>P. aeruginosa</i> (95.3%)
Purity (post-freeze) 7 days at 37°C in an aerobic atmosphere with and without 5%CO ₂ on Tryptic Soy agar with 5% defibrinated sheep blood	Growth consistent with expected colony morphology	Growth consistent with expected colony morphology
Viability (post-freeze)	Growth	Growth

¹Relatedness between bacterial strains has traditionally been determined using DDH. For additional information, refer to Auch, A. F., et al. "Digital DNA-DNA Hybridization for Microbial Species Delineation by Means of Genome-to-Genome Sequence Comparison." *Stand. Genomic Sci.* 2 (2010): 117-134. PubMed: 21304684.

Table 3: *Pseudomonas aeruginosa*, Strain NB52019-LpxC_{G208S} (NR-51886)

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis Cellular morphology Colony morphology Motility (wet mount) VITEK® MS (MALDI-TOF)	Gram-negative rods Report results Report results <i>P. aeruginosa</i>	Gram-negative rods Circular, convex, entire, smooth and cream Motile <i>P. aeruginosa</i> (99.9%)
Antibiotic Susceptibility Profile Etest® antibiotic test strips 1 day at 35°C in an aerobic atmosphere on Mueller Hinton agar Chloramphenicol Ofloxacin	Report results Report results	32 µg/mL 1.5 µg/mL
Genotypic Analysis Confirmation of mutations Digital DNA-DNA hybridization (dDDH) ¹	Mutations confirmed ≥ 70% for species identification	Pending <i>P. aeruginosa</i> (95.3%)
Purity (post-freeze) 7 days at 37°C in an aerobic atmosphere with and without 5% CO ₂ on Tryptic Soy agar with 5% defibrinated sheep blood	Growth consistent with expected colony morphology	Growth consistent with expected colony morphology
Viability (post-freeze)	Growth	Growth

¹Relatedness between bacterial strains has traditionally been determined using DDH. For additional information, refer to Auch, A. F., et al. "Digital DNA-DNA Hybridization for Microbial Species Delineation by Means of Genome-to-Genome Sequence Comparison." *Stand. Genomic Sci.* 2 (2010): 117-134. PubMed: 21304684.

Table 4: *Pseudomonas aeruginosa*, Strain NB52019-CDJ0037 (NR-51887)

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis Cellular morphology Colony morphology Motility (wet mount) VITEK® MS (MALDI-TOF)	Gram-negative rods Report results Report results <i>P. aeruginosa</i>	Gram-negative rods Irregular, low convex, undulate, rough and cream Motile <i>P. aeruginosa</i> (99.9%)
Antibiotic Susceptibility Profile Etest® antibiotic test strips 1 day at 35°C in an aerobic atmosphere on Mueller Hinton agar Chloramphenicol Ofloxacin	Report results Report results	64 µg/mL 1.0 µg/mL
Genotypic Analysis Confirmation of mutations Digital DNA-DNA hybridization (dDDH) ¹	Mutations confirmed ≥ 70% for species identification	Pending <i>P. aeruginosa</i> (95.2%)
Purity (post-freeze) 7 days at 37°C in an aerobic atmosphere with and without 5% CO ₂ on Tryptic Soy agar with 5% defibrinated sheep blood	Growth consistent with expected colony morphology	Growth consistent with expected colony morphology
Viability (post-freeze)	Growth	Growth

¹Relatedness between bacterial strains has traditionally been determined using DDH. For additional information, refer to Auch, A. F., et al. "Digital DNA-DNA Hybridization for Microbial Species Delineation by Means of Genome-to-Genome Sequence Comparison." *Stand. Genomic Sci.* 2 (2010): 117-134. PubMed: 21304684.

Table 5: *Pseudomonas aeruginosa*, Strain NB52217-P2 (NR-51888)

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis Cellular morphology Colony morphology Motility (wet mount) VITEK® MS (MALDI-TOF)	Gram-negative rods Report results Report results <i>P. aeruginosa</i>	Gram-negative rods Circular, convex, entire, smooth and cream Motile <i>P. aeruginosa</i> (99.9%)
Antibiotic Susceptibility Profile Etest® antibiotic test strips 1 day at 35°C in an aerobic atmosphere on Mueller Hinton agar Chloramphenicol Ofloxacin	Report results Report results	256 µg/mL 3 µg/mL
Genotypic Analysis Confirmation of mutations Digital DNA-DNA hybridization (dDDH) ¹	Mutations confirmed ≥ 70% for species identification	Pending <i>P. aeruginosa</i> (95.3%)
Purity (post-freeze) 7 days at 37°C in an aerobic atmosphere with 5% CO ₂ on Tryptic Soy agar with 5% defibrinated sheep blood	Growth consistent with expected colony morphology	Growth consistent with expected colony morphology
Viability (post-freeze)	Growth	Growth

¹Relatedness between bacterial strains has traditionally been determined using DDH. For additional information, refer to Auch, A. F., et al. "Digital DNA-DNA Hybridization for Microbial Species Delineation by Means of Genome-to-Genome Sequence Comparison." *Stand. Genomic Sci.* 2 (2010): 117-134. PubMed: 21304684.

Table 6: *Pseudomonas aeruginosa*, Strain NB52217-P6 (NR-51889)

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis Cellular morphology Colony morphology Motility (wet mount) VITEK® MS (MALDI-TOF)	Gram-negative rods Report results Report results <i>P. aeruginosa</i>	Gram-negative rods Circular, convex, entire, smooth and cream Motile <i>P. aeruginosa</i> (99.9%)
Antibiotic Susceptibility Profile Etest® antibiotic test strips 1 day at 35°C in an aerobic atmosphere on Mueller Hinton agar Chloramphenicol Ofloxacin	Report results Report results	≥ 256 µg/mL 12 to 16 µg/mL
Genotypic Analysis Confirmation of mutations Digital DNA-DNA hybridization (dDDH) ¹	Mutations confirmed ≥ 70% for species identification	Pending <i>P. aeruginosa</i> (95.3%)
Purity (post-freeze) 7 days at 37°C in an aerobic atmosphere with and without 5% CO ₂ on Tryptic Soy agar	Growth consistent with expected colony morphology	Growth consistent with expected colony morphology
Viability (post-freeze)	Growth	Growth

¹Relatedness between bacterial strains has traditionally been determined using DDH. For additional information, refer to Auch, A. F., et al. "Digital DNA-DNA Hybridization for Microbial Species Delineation by Means of Genome-to-Genome Sequence Comparison." *Stand. Genomic Sci.* 2 (2010): 117-134. PubMed: 21304684.

Table 7: *Pseudomonas aeruginosa*, Strain NB52217-P13 (NR-51890)

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis Cellular morphology Colony morphology Motility (wet mount) VITEK® MS (MALDI-TOF)	Gram-negative rods Report results Report results <i>P. aeruginosa</i>	Gram-negative rods Circular, low convex, entire, smooth and cream Motile <i>P. aeruginosa</i> (99.9%)
Antibiotic Susceptibility Profile Etest® antibiotic test strips 1 day at 35°C in an aerobic atmosphere on Mueller Hinton agar Chloramphenicol Ofloxacin	Report results Report results	> 256 µg/mL > 32 µg/mL
Genotypic Analysis Confirmation of mutations Digital DNA-DNA hybridization (dDDH) ¹	Mutations confirmed ≥ 70% for species identification	Pending <i>P. aeruginosa</i> (95.3%)
Purity (post-freeze) 7 days at 37°C in an aerobic atmosphere with and without 5% CO ₂ on Tryptic Soy agar with 5% defibrinated sheep blood	Growth consistent with expected colony morphology	Growth consistent with expected colony morphology
Viability (post-freeze)	Growth	Growth

¹Relatedness between bacterial strains has traditionally been determined using DDH. For additional information, refer to Auch, A. F., et al. "Digital DNA-DNA Hybridization for Microbial Species Delineation by Means of Genome-to-Genome Sequence Comparison." *Stand. Genomic Sci.* 2 (2010): 117-134. PubMed: 21304684.

Table 8: *Pseudomonas aeruginosa*, Strain NB52217-PA4465_{N193T} (NR-51891)

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis		
Cellular morphology Colony morphology	Gram-negative rods Report results	Gram-negative rods Circular, convex, entire, smooth and cream
Motility (wet mount) VITEK® MS (MALDI-TOF)	Report results <i>P. aeruginosa</i>	Motile <i>P. aeruginosa</i> (99.9%)
Antibiotic Susceptibility Profile		
Etest® antibiotic test strips 1 day at 35°C in an aerobic atmosphere on Mueller Hinton agar Chloramphenicol Ofloxacin	Report results Report results	128 µg/mL 2 µg/mL
Genotypic Analysis		
Confirmation of mutations Digital DNA-DNA hybridization (dDDH) ¹	Mutations confirmed ≥ 70% for species identification	Pending <i>P. aeruginosa</i> (95.3%)
Purity (post-freeze)		
7 days at 37°C in an aerobic atmosphere with and without 5% CO ₂ on Tryptic Soy agar with 5% defibrinated sheep blood	Growth consistent with expected colony morphology	Growth consistent with expected colony morphology
Viability (post-freeze)		
	Growth	Growth

¹Relatedness between bacterial strains has traditionally been determined using DDH. For additional information, refer to Auch, A. F., et al. "Digital DNA-DNA Hybridization for Microbial Species Delineation by Means of Genome-to-Genome Sequence Comparison." *Stand. Genomic Sci.* 2 (2010): 117-134. PubMed: 21304684.

Table 9: *Pseudomonas aeruginosa*, Strain NB52200-P6a (NC) (NR-51892)

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis		
Cellular morphology Colony morphology	Gram-negative rods Report results	Gram-negative rods Circular, convex, entire, smooth and cream
Motility (wet mount) VITEK® MS (MALDI-TOF)	Report results <i>P. aeruginosa</i>	Motile <i>P. aeruginosa</i> (99.9%)
Antibiotic Susceptibility Profile		
Etest® antibiotic test strips 1 day at 35°C in an aerobic atmosphere on Mueller Hinton agar Chloramphenicol Ofloxacin	Report results Report results	2 µg/mL 0.064 µg/mL
Genotypic Analysis		
Confirmation of mutations Digital DNA-DNA hybridization (dDDH) ¹	Mutations confirmed ≥ 70% for species identification	Pending <i>P. aeruginosa</i> (95.4%)
Purity (post-freeze)		
7 days at 37°C in an aerobic atmosphere with and without 5%CO ₂ on Tryptic Soy agar	Growth consistent with expected colony morphology	Growth consistent with expected colony morphology
Viability (post-freeze)		
	Growth	Growth

¹Relatedness between bacterial strains has traditionally been determined using DDH. For additional information, refer to Auch, A. F., et al. "Digital DNA-DNA Hybridization for Microbial Species Delineation by Means of Genome-to-Genome Sequence Comparison." *Stand. Genomic Sci.* 2 (2010): 117-134. PubMed: 21304684.

Table 10: *Pseudomonas aeruginosa*, Strain NB52200-P6b (SC) (NR-51893)

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis Cellular morphology Colony morphology Motility (wet mount) VITEK® MS (MALDI-TOF)	Gram-negative rods Report results Report results <i>P. aeruginosa</i>	Gram-negative rods Circular, convex, entire, smooth and cream Motile <i>P. aeruginosa</i> (99.9%)
Antibiotic Susceptibility Profile Etest® antibiotic test strips 1 day at 35°C in an aerobic atmosphere on Mueller Hinton agar Chloramphenicol Ofloxacin	Report results Report results	1.5 to 2 µg/mL 0.047 µg/mL
Genotypic Analysis Confirmation of mutations Digital DNA-DNA hybridization (dDDH) ¹	Mutations confirmed ≥ 70% for species identification	Pending <i>P. aeruginosa</i> (95.3%)
Purity (post-freeze) 7 days at 37°C in an aerobic atmosphere with and without 5% CO ₂ on Tryptic Soy agar	Growth consistent with expected colony morphology	Growth consistent with expected colony morphology
Viability (post-freeze)	Growth	Growth

¹Relatedness between bacterial strains has traditionally been determined using DDH. For additional information, refer to Auch, A. F., et al. "Digital DNA-DNA Hybridization for Microbial Species Delineation by Means of Genome-to-Genome Sequence Comparison." *Stand. Genomic Sci.* 2 (2010): 117-134. PubMed: 21304684.

Table 11: *Pseudomonas aeruginosa*, Strain NB52200-P13a (NC) (NR-51894)

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis Cellular morphology Colony morphology Motility (wet mount) VITEK® MS (MALDI-TOF)	Gram-negative rods Report results Report results <i>P. aeruginosa</i>	Gram-negative rods Circular, low convex, entire, smooth and cream Motile <i>P. aeruginosa</i> (99.9%)
Antibiotic Susceptibility Profile Etest® antibiotic test strips 1 day at 35°C in an aerobic atmosphere on Mueller Hinton agar Chloramphenicol Ofloxacin	Report results Report results	8 µg/mL 0.047 µg/mL
Genotypic Analysis Confirmation of mutations Digital DNA-DNA hybridization (dDDH) ¹	Mutations confirmed ≥ 70% for species identification	Pending <i>P. aeruginosa</i> (95.3%)
Purity (post-freeze) 7 days at 37°C in an aerobic atmosphere with and without 5%CO ₂ on Tryptic Soy agar	Growth consistent with expected colony morphology	Growth consistent with expected colony morphology
Viability (post-freeze)	Growth	Growth

¹Relatedness between bacterial strains has traditionally been determined using DDH. For additional information, refer to Auch, A. F., et al. "Digital DNA-DNA Hybridization for Microbial Species Delineation by Means of Genome-to-Genome Sequence Comparison." *Stand. Genomic Sci.* 2 (2010): 117-134. PubMed: 21304684.

Table 12: *Pseudomonas aeruginosa*, Strain NB52200-P13b (SC) (NR-51895)

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis Cellular morphology Colony morphology Motility (wet mount) VITEK® MS (MALDI-TOF)	Gram-negative rods Report results Report results <i>P. aeruginosa</i>	Gram-negative rods Circular, convex, entire, smooth and cream Motile <i>P. aeruginosa</i> (99.9%)
Antibiotic Susceptibility Profile Etest® antibiotic test strips 1 day at 35°C in an aerobic atmosphere on Mueller Hinton agar Chloramphenicol Ofloxacin	Report results Report results	2 to 3 µg/mL 0.006 to 0.008 µg/mL
Genotypic Analysis Confirmation of mutations Digital DNA-DNA hybridization (dDDH) ¹	Mutations confirmed ≥ 70% for species identification	Pending <i>P. aeruginosa</i> (95.3%)
Purity (post-freeze) 7 days at 37°C in an aerobic atmosphere with and without 5%CO ₂ on Tryptic Soy agar	Growth consistent with expected colony morphology	Growth consistent with expected colony morphology
Viability (post-freeze)	Growth	Growth

¹Relatedness between bacterial strains has traditionally been determined using DDH. For additional information, refer to Auch, A. F., et al. "Digital DNA-DNA Hybridization for Microbial Species Delineation by Means of Genome-to-Genome Sequence Comparison." *Stand. Genomic Sci.* 2 (2010): 117-134. PubMed: 21304684.

Table 13: *Pseudomonas aeruginosa*, Strain NB52042-CDJ0042 (NR-51896)

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis Cellular morphology Colony morphology Motility (wet mount) VITEK® MS (MALDI-TOF)	Gram-negative rods Report results Report results <i>P. aeruginosa</i>	Gram-negative rods Irregular, low convex, undulate, smooth and cream Motile <i>P. aeruginosa</i> (99.9%)
Antibiotic Susceptibility Profile Etest® antibiotic test strips 1 day at 35°C in an aerobic atmosphere on Mueller Hinton agar Chloramphenicol Ofloxacin	Report results Report results	48 to 64 µg/mL 1.5 to 2 µg/mL
Genotypic Analysis Confirmation of mutations Digital DNA-DNA hybridization (dDDH) ¹	Mutations confirmed ≥ 70% for species identification	Pending <i>P. aeruginosa</i> (95.3%)
Purity (post-freeze) 7 days at 37°C in an aerobic atmosphere on Tryptic Soy agar	Growth consistent with expected colony morphology	Growth consistent with expected colony morphology
7 days at 37°C in an aerobic atmosphere with 5%CO ₂ on Tryptic Soy agar with 5% defibrinated sheep blood	Growth consistent with expected colony morphology	Growth consistent with expected colony morphology
Viability (post-freeze)	Growth	Growth

¹Relatedness between bacterial strains has traditionally been determined using DDH. For additional information, refer to Auch, A. F., et al. "Digital DNA-DNA Hybridization for Microbial Species Delineation by Means of Genome-to-Genome Sequence Comparison." *Stand. Genomic Sci.* 2 (2010): 117-134. PubMed: 21304684.

Table 14: *Pseudomonas aeruginosa*, Strain NB52019-CDR0026 (NR-51898)

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis Cellular morphology Colony morphology Motility (wet mount) VITEK® MS (MALDI-TOF)	Gram-negative rods Report results Report results <i>P. aeruginosa</i>	Gram-negative rods Irregular, low convex, undulate, rough and cream Motile <i>P. aeruginosa</i> (99.9%)
Antibiotic Susceptibility Profile Etest® antibiotic test strips 1 day at 35°C in an aerobic atmosphere on Mueller Hinton agar Chloramphenicol Ofloxacin	Report results Report results	96 µg/mL 0.75 µg/mL
Genotypic Analysis Confirmation of mutations Digital DNA-DNA hybridization (dDDH) ¹	Mutations confirmed ≥ 70% for species identification	Pending <i>P. aeruginosa</i> (95.3%)
Purity (post-freeze) 7 days at 37°C in an aerobic atmosphere with and without 5% CO ₂ on Tryptic Soy agar with 5% defibrinated sheep blood	Growth consistent with expected colony morphology	Growth consistent with expected colony morphology
Viability (post-freeze)	Growth	Growth

¹Relatedness between bacterial strains has traditionally been determined using DDH. For additional information, refer to Auch, A. F., et al. "Digital DNA-DNA Hybridization for Microbial Species Delineation by Means of Genome-to-Genome Sequence Comparison." *Stand. Genomic Sci.* 2 (2010): 117-134. PubMed: 21304684.

Table 15: *Pseudomonas aeruginosa*, Strain NB52019-CDR0061 (NR-51899)

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis Cellular morphology Colony morphology Motility (wet mount) VITEK® MS (MALDI-TOF)	Gram-negative rods Report results Report results <i>P. aeruginosa</i>	Gram-negative rods Irregular, convex, undulate, rough and cream Motile <i>P. aeruginosa</i> (99.9%)
Antibiotic Susceptibility Profile Etest® antibiotic test strips 1 day at 35°C in an aerobic atmosphere on Mueller Hinton agar Chloramphenicol Ofloxacin	Report results Report results	> 256 µg/mL 1 µg/mL
Genotypic Analysis Confirmation of mutations Digital DNA-DNA hybridization (dDDH) ¹	Mutations confirmed ≥ 70% for species identification	Pending <i>P. aeruginosa</i> (95.3%)
Purity (post-freeze) 7 days at 37°C in an aerobic atmosphere with and without 5% CO ₂ on Tryptic Soy agar with 5% defibrinated sheep blood	Growth consistent with expected colony morphology	Growth consistent with expected colony morphology
Viability (post-freeze)	Growth	Growth

¹Relatedness between bacterial strains has traditionally been determined using DDH. For additional information, refer to Auch, A. F., et al. "Digital DNA-DNA Hybridization for Microbial Species Delineation by Means of Genome-to-Genome Sequence Comparison." *Stand. Genomic Sci.* 2 (2010): 117-134. PubMed: 21304684.

Table 16: *Pseudomonas aeruginosa*, Strain NB52019-CDJ0011 (NR-51900)

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis Cellular morphology Colony morphology Motility (wet mount) VITEK® MS (MALDI-TOF)	Gram-negative rods Report results Report results <i>P. aeruginosa</i>	Gram-negative rods Circular, low convex, undulate, smooth and cream Motile <i>P. aeruginosa</i> (99.9%)
Antibiotic Susceptibility Profile Etest® antibiotic test strips 1 day at 35°C in an aerobic atmosphere on Mueller Hinton agar Chloramphenicol Ofloxacin	Report results Report results	48 µg/mL 1.5 µg/mL
Genotypic Analysis Confirmation of mutations Digital DNA-DNA hybridization (dDDH) ¹	Mutations confirmed ≥ 70% for species identification	Pending <i>P. aeruginosa</i> (95.3%)
Purity (post-freeze) 7 days at 37°C in an aerobic atmosphere with and without 5% CO ₂ on Tryptic Soy agar with 5% defibrinated sheep blood	Growth consistent with expected colony morphology	Growth consistent with expected colony morphology
Viability (post-freeze)	Growth	Growth

¹Relatedness between bacterial strains has traditionally been determined using DDH. For additional information, refer to Auch, A. F., et al. "Digital DNA-DNA Hybridization for Microbial Species Delineation by Means of Genome-to-Genome Sequence Comparison." *Stand. Genomic Sci.* 2 (2010): 117-134. PubMed: 21304684.

Table 17: *Pseudomonas aeruginosa*, Strain NB52203-CDB0011 (NR-51902)

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis Cellular morphology Colony morphology Motility (wet mount) VITEK® MS (MALDI-TOF)	Gram-negative rods Report results Report results <i>P. aeruginosa</i>	Gram-negative rods Irregular, low convex, undulate, smooth and cream Motile <i>P. aeruginosa</i> (99.9%)
Antibiotic Susceptibility Profile Etest® antibiotic test strips 1 day at 35°C in an aerobic atmosphere on Mueller Hinton agar Chloramphenicol Ofloxacin	Report results Report results	> 256 µg/mL 1.5 µg/mL
Genotypic Analysis Confirmation of mutations Digital DNA-DNA hybridization (dDDH) ¹	Mutations confirmed ≥ 70% for species identification	Pending <i>P. aeruginosa</i> (90.5%)
Purity (post-freeze) 7 days at 37°C in an aerobic atmosphere with and without 5% CO ₂ on Tryptic Soy agar	Growth consistent with expected colony morphology	Growth consistent with expected colony morphology
Viability (post-freeze)	Growth	Growth

¹Relatedness between bacterial strains has traditionally been determined using DDH. For additional information, refer to Auch, A. F., et al. "Digital DNA-DNA Hybridization for Microbial Species Delineation by Means of Genome-to-Genome Sequence Comparison." *Stand. Genomic Sci.* 2 (2010): 117-134. PubMed: 21304684.

/Sonia Bjorum Brower/

Sonia Bjorum Brower

28 SEP 2023

Technical Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected by ATCC® and the contributor to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

